



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Then $us = a\theta \dots (1)$. $\therefore s = \frac{a\theta}{u} = \frac{am\theta}{m}$ is the intrinsic equation to the curve.

$$\text{From (1) } \frac{ds}{d\theta} = \frac{a}{u} = \sqrt{r^2 + \left(\frac{dr}{d\theta}\right)^2}.$$

$$\therefore r^2 + \left(\frac{dr}{d\theta}\right)^2 = \frac{a^2}{u^2} \text{ and } d\theta = \frac{dr}{\sqrt{\left(\frac{a^2}{u^2} - r^2\right)}}.$$

$$\therefore \theta = \sin^{-1} \frac{ur}{a} \dots (2).$$

$\therefore r = \frac{a}{u} \sin\theta = \frac{am}{m} \sin\theta$, is the polar equation
and $m(x^2 + y^2) = any$, is the rectangular equation.

The value of θ from (2) in (1) gives

$$s = \frac{a}{u} \sin^{-1} \frac{ur}{a},$$

the length for any value of r . When $r = a$,

$$s = \frac{a}{u} \sin^{-1} u = \frac{am}{m} \sin^{-1} \frac{m}{n} = \text{distance run.}$$

$$t = \text{time} = \frac{s}{n} = \frac{a}{m} \sin^{-1} \frac{m}{n}.$$

Also solved by Professors O. W. Anthony, J. Scheffer, and William Symmonds.

28. Proposed by F. P. MATZ, M. Sc., Ph. D., Professor of Mathematics and Astronomy in New Windsor College, New Windsor, Maryland.

How far from the stage must Miss Love sit in order that she may see to best advantage Mr. Rich deliver the valedictory oration?

Solution by O. W. ANTHONY, Missouri Military Academy, Mexico, Missouri, and the PROPOSER.

Let E represent the position of Miss Love's eyes; DB the stage from which Mr. Rich orates; AB , $=m$ feet, the height of the stage above Miss Love's eyes; BC , $=n$ feet, the height of Mr. Rich; and AE , $=x$ feet, the required distance. In order that Miss Love may see Mr. Rich to best advantage, the angle BEC must be a maximum, that is,



$$U = \tan^{-1} \left(\frac{m+n}{x} \right) - \tan^{-1} \left(\frac{m}{x} \right) = \text{a Maximum.}$$

$$\therefore \frac{dU}{dx} = \frac{m}{x^2 + m^2} - \frac{m+n}{x^2 + (m+n)^2} = 0 \dots (1).$$

Whence $x = \sqrt{m(m+n)}$ feet, which is the required distance.

29. Proposed by CHARLES E. MYERS, Canton, Ohio

A hen running at the rate of $n=2$ feet per second, on the circumference of a circle, radius $r=50$ feet, is observed by a hawk $a=600$ feet directly above the center.



The hawk at once starts in pursuit, flying at the rate of $m=5$ feet per second and keeping always in a straight line with the starting point and the hen.

Determine the path followed and the distance the hawk will fly before catching the hen.

Solution by G. B. M. ZERR, A. M., Principal of High School, Staunton, Virginia.

Let the origin be at the vertex of the cone around which the path of the hawk winds. σ =length of the hawk's path, s =the length of the projection of this path on the plane (x,y) , ρ =radius vector of this projection,

$\frac{n}{m}=u$, $\frac{r}{a}=c$. Then $x^2+y^2=c^2z^2$ is the equation of the cone, also $n\sigma=r\theta$,

where θ is the angle subtended by the hen's path at the centre of the circle.

$$\therefore d\sigma = \frac{r}{u} d\theta = \sqrt{dx^2 + dy^2 + dz^2} = \sqrt{ds^2 + dz^2}$$

$$= \left\{ \rho^2 + \left(\frac{d\rho}{d\theta} \right)^2 + \left(\frac{dz}{d\theta} \right)^2 \right\}^{\frac{1}{2}} d\theta, \text{ but } \rho^2 = x^2 + y^2 = c^2 z^2, \therefore \rho = cz, \therefore dz = \frac{d\rho}{c}.$$

$$\therefore \frac{d\sigma}{d\theta} = \frac{r}{u} = \left\{ \rho^2 + \left(\frac{d\rho}{d\theta} \right)^2 + \frac{1}{c^2} \left(\frac{d\rho}{d\theta} \right)^2 \right\}^{\frac{1}{2}}.$$

$$\therefore \frac{d\theta}{d\rho} = \frac{\sqrt{c^2+1}}{c} \cdot \frac{1}{\left\{ \frac{\rho^2}{u^2} - \rho^2 \right\}^{\frac{1}{2}}}, \quad \theta = \frac{\sqrt{c^2+1}}{c} \sin^{-1} \frac{n\rho}{r}.$$

$$\therefore \theta = \frac{\sqrt{r^2+a^2}}{r} \sin^{-1} \left(\frac{n\rho}{mr} \right); \quad \rho = \frac{mr}{n} \sin \frac{r\theta}{\sqrt{r^2+a^2}}.$$

$$z = \rho / c = \frac{mn}{n} \sin \frac{r\theta}{\sqrt{r^2+a^2}} = 1500 \sin \frac{\theta}{\sqrt{145}}.$$

$$x = \rho \cos \theta = \frac{mr}{n} \sin \frac{r\theta}{\sqrt{r^2+a^2}} \cos \theta = 125 \sin \frac{\theta}{\sqrt{145}} \cos \theta.$$

$$y = \rho \sin \theta = \frac{mr}{n} \sin \frac{r\theta}{\sqrt{r^2+a^2}} \sin \theta = 125 \sin \frac{\theta}{\sqrt{145}} \sin \theta.$$

These values of x , y , z determine the hawk's path.

$$\text{Also } n\sigma = r\theta, \therefore \sigma = \frac{mr}{n} \left[\frac{\sqrt{r^2+a^2}}{r} \sin^{-1} \left(\frac{n\rho}{mr} \right) \right]_0^r = \frac{m\sqrt{r^2+a^2}}{u} \sin^{-1} \left(\frac{n}{m} \right).$$

$$\therefore \sigma = 125 \sqrt{145} \sin^{-1} \left(\frac{2}{5} \right) = 619.406 \text{ feet approximately, the distance}$$

the hawk flies before catching the hen.

Solved in a similar manner by Professor J. F. W. Scheffer.